

Name \_\_\_\_\_

## Solution Concentration (Molarity) and Dilution

- 1) What is the molarity of a solution if 58.2g of sodium chloride was dissolved in sufficient water to prepare a 1.83L solution?
- 2) How many grams of sodium hydroxide must be dissolved to produce 600.mL of a 0.750M solution?
- 3) If 9.62g of potassium iodide is dissolved in sufficient water to make a 4.50L solution, then what is its molarity?
- 4) What volume of solution do you have if you know you dissolved 0.0500g nitric acid to make a solution that was determined to be 0.0793M?
- 5) If 6.32g of barium sulfate is produced from the reaction of sulfuric acid with barium chloride, then what was the molarity of the sulfuric acid solution if only 50.0mL was used?
- 6) If 98.2mL of a 1.50M hydrochloric acid solution is reacted with excess aluminum hydroxide, then how many grams of aluminum chloride would be your theoretical yield?  
$$\text{Al(OH)}_3 + 3\text{HCl} \rightarrow \text{AlCl}_3 + 3\text{H}_2\text{O}$$
- 7) If 25.2mL of a 2.50M sodium bromide solution reacts with excess chlorine gas, then how many liters of bromine gas would you expect to collect at 1.03atm and 295K?
- 8) What was the initial concentration of a silver nitrate solution if 45.8mL of the solution was reacted with magnesium, and 3.28g of silver was recovered?
- 9) How many milliliters of a 1.50M acetic acid solution will be needed to create 2.50L of a 0.250M solution?
- 10) Concentrated hydrochloric acid has a concentration of 12.1 moles per liter. How many milliliters of the concentrated acid should be used to create 3.25L of a 0.100M solution?
- 11) If you want to produce 1.00L of 0.050M nitric acid from a 10.0M solution, then calculate the volume of nitric acid and the volume of water necessary to make the solution.